

UNCLASSIFIED

AD NUMBER
AD810432
NEW LIMITATION CHANGE
TO Approved for public release, distribution unlimited
FROM Distribution authorized to U.S. Gov't. agencies and their contractors; Administrative/Operational Use; MAR 1967. Other requests shall be referred to Department of the Army, Fort Detrick, Attn: Technical Releases Branch, Frederick, MD 21701.
AUTHORITY
Army Biological Defense Research Lab. ltr dtd 28 Sep 1971

THIS PAGE IS UNCLASSIFIED

AD

810432

TECHNICAL MANUSCRIPT 369

PROTECTIVE ACTION OF PROTEIN
ON NEWLY GERMINATED
BACILLUS ANTHRACIS SPORES IN TISSUE BREI

William I. Jones, Jr.


Ralph E. Lincoln

Byron U. Ross

MARCH 1967

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

DISCLAIMER NOTICE



**THIS DOCUMENT IS BEST
QUALITY AVAILABLE. THE COPY
FURNISHED TO DTIC CONTAINED
A SIGNIFICANT NUMBER OF
PAGES WHICH DO NOT
REPRODUCE LEGIBLY.**

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland 21701

TECHNICAL MANUSCRIPT 369

PROTECTIVE ACTION OF PROTEIN ON NEWLY GERMINATED
BACILLUS ANTHRACIS SPORES IN TISSUE BREI

William I. Jones, Jr.

Ralph E. Lincoln

Byron U. Ross

Process Development Division
AGENT DEVELOPMENT AND ENGINEERING LABORATORY

Project IC522301A059

March 1967

In conducting the research described in this report, the investigators adhered to the "Guide for Laboratory Animal Facilities and Care," as promulgated by the Committee on the Guide for Laboratory Animal Facilities and Care of the Institute of Laboratory Animal Resources, National Academy of Sciences-National Research Council.

PROTECTIVE ACTION OF PROTEIN ON NEWLY GERMINATED
BACILLUS ANTHRACIS SPORES IN TISSUE BREI

ABSTRACT

A complete loss of viability was observed when newly germinated spores of Bacillus anthracis were placed in contact with freshly ground mouse tissue. This loss was overcome by the addition of 0.18% Bacto beef extract and 0.3% Bacto peptone to the amino acid germinating mixture.

While developing procedures for determining in vivo growth curves of Bacillus anthracis,¹ we observed that the conditions of germination of the spores markedly influenced the initial destruction of cells and our ability to account for the dose quantitatively.

Howie and Cruickshank² found viable anthrax spores in the spleen of mice killed 3 months after inoculation with B. anthracis spores. In order that dormant spores not be present in the host to affect the later quantitation of in vivo growth, a challenge with germinated spores was chosen. Heat-shocked spores were germinated in a medium containing 0.0125% L-alanine, 0.0125% L-tyrosine, and 0.00625% adenosine by placing 1×10^8 heat-shocked spores contained in one ml into a test tube containing 3 ml of the stock amino acid mixture and 6 ml of sterile distilled water.³ The resulting suspension was incubated at 37 C for 30 minutes. Vegetative growth occurs if the spores are transferred to a suitable growth medium. The condition of the organism, i.e., dormant spore, germinated spore, or vegetative organism, was determined by phase microscopy and differential staining. The percentage germination was determined by differential plate count before and after the addition of sufficient phenol to obtain a final concentration of 1%.⁴

For this work, we believed it essential to be able to account, immediately after their introduction into the host, for 100% of the organisms used as inoculum. The steps following inoculation were to immediately sacrifice the animal, skin, grind in a Waring Blendor, then quantitatively dilute in gelatin phosphate buffer (pH 7.2) and plate onto nutrient agar. When spores were germinated in the amino acid mixture, no organisms could be recovered when assayed as described. To determine whether this observation was attributable to physical destruction of the organisms by the grinding mechanism or to some anthracidal material of normal tissue, germinated

spores were added to a suspension of uninfected (control) tissue and a viable plate count was made. Once again there was a complete loss of organisms. This finding was in complete agreement with that of Bloom et al.⁶

When 0.18% Bacto beef extract and 0.3% Bacto peptone were added to the amino acid mixture, the germinated spores were completely protected from the germicidal action of ground tissue or of tissue during grinding. Thus, we were able to account for and recover 100% of the organisms injected into a live animal.

This change of resistance in germinated spores may be due to a more advanced stage of development occurring in the presence of protein. As observed by light microscopy, there is no real difference in rate or percentage of spores losing refractivity in the two mixtures. Spores left in the amino acid mixture for 90 minutes will not develop into vegetative cells, whereas about 80 to 85% of the spores have divided in the protein-containing mixture. Because both dormant spores and newly germinated spores respond similarly, we suggest that resistance cells have developed the Embden-Meyerhof enzyme system, which is non-operative in dormant spores or in the initial stages of germinated spores.⁶ However, other explanations are possible; for example, sensitivity to lysozyme may change in the presence of spermine.⁷

These data show that the newly germinated spore is very sensitive to destruction in the animal body. This fact needs to be considered in views on dynamics of infection, particularly in regard to host resistance and initiation of disease by different routes.

LITERATURE CITED

1. Jones, W.I., Jr.; Klein, F.; Lincoln, R.E.; Walker, J.S.; Mahlandt, B.G.; Dobbs, J.P. February 1967. Anthrax infection in mice as a model for studying the dynamics of therapy in bacterial diseases, (Technical Manuscript 347). Process Development Division, Fort Detrick, Frederick, Maryland.
2. Howie, J.W.; Cruickshank, J. 1947. Effect of shock-producing substances on experimental anthrax infection in mice. J. Pathol. Bacteriol. 59:127-135.
3. Hills, G.M. 1949. Chemical factors in the germination of spore-bearing aerobes: The effects of amino-acids on the germination of Bacillus anthracis with some observations on the relation of optical form to biological activity. Biochem. J. 45:363-370.
4. Fernelius, A.L. 1960. Comparison of two heat-shock methods and a phenol treatment method for determining germination rates of spores of Bacillus anthracis. J. Bacteriol. 79:755-756.
5. Bloom, W.L.; Watson, D.W.; Cromartie, W.J.; Freed, M. 1947. Studies on infection with Bacillus anthracis: IV. Preparation and characterization of an anthracidal substance from various animal tissues. J. Infect. Dis. 60:41-52.
6. Blumenthal, H.J. 1965. Changes in carbohydrate metabolism of Bacillus cereus spores during the return to vegetative growth, p. 222-236. In H.O. Halvorsen (ed.) Spores III. Burgess Publishing Company, Minneapolis, Minn.
7. Wright, G.G. 1961. Influence of spermine and related substances on susceptibility of Bacillus anthracis to lysozyme. Proc. Soc. Exp. Biol. Med. 108:740-742.

Unclassified
Security Classification

DOCUMENT CONTROL DATA - R&D		
(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)		
1 ORIGINATING ACTIVITY (Corporate author) Department of the Army Fort Detrick, Frederick, Maryland 21701		2a REPORT SECURITY CLASSIFICATION Unclassified 2b GROUP
3 REPORT TITLE PROTECTIVE ACTION OF PROTEIN ON NEWLY GERMINATED <u>BACILLUS ANTHRACIS</u> SPORES IN TISSUE BREI		
4 DESCRIPTIVE NOTES (Type of report and inclusive dates)		
5 AUTHOR(S) (Last name, first name, initial) Jones, William I., Jr. Ross, Byron U. Lincoln, Ralph E.		
6 REPORT DATE March 1967	7a TOTAL NO OF PAGES 8	7b NO OF REFS 7
8a CONTRACT OR GRANT NO. b PROJECT NO. 1C522301A059 c. d.	9a ORIGINATOR'S REPORT NUMBER(S) Technical Manuscript 369 9b OTHER REPORT NO(S) (Any other numbers that may be assigned this report)	
10 AVAILABILITY/LIMITATION NOTICES Qualified requesters may obtain copies of this publication from DDC. Foreign announcement and dissemination of this publication by DDC is not authorized. Release or announcement to the public is not authorized.		
11 SUPPLEMENTARY NOTES	12 SPONSORING MILITARY ACTIVITY Department of the Army Fort Detrick, Frederick, Maryland 21701	
13 ABSTRACT A complete loss of viability was observed when newly germinated spores of <u>Bacillus anthracis</u> were placed in contact with freshly ground mouse tissue. This loss was overcome by the addition of 0.18% Bacto beef extract and 0.3% Bacto peptone to the amino acid germinating mixture.		
14. Key Words * <u>Bacillus anthracis</u> Proteins *Spores Culture media *Germination Tissues Amino acids		

DD FORM 1473
1 JAN 64

Unclassified
Security Classification